



Comparison of Loose Smut, Common Bunt, and Karnal Bunt of Wheat

Bob Hunger (Extension Wheat Pathologist), Brian Olson (Extension Assistant & Plant Disease Diagnostician), & Larry Singleton (Professor of Plant Pathology.)
Department of Entomology and Plant Pathology, Oklahoma State University




July 2001

The bunts and smuts are caused by fungi that affect the seed of wheat and other small grains and grasses. Only loose smut (Fig. 1) and common bunt (Fig. 2 - also called stinking smut) occur in Oklahoma. However, Karnal bunt (Fig. 3 - also called partial bunt), which does not occur in Oklahoma, is a concern because grain that tests positive for Karnal bunt can not move freely into international markets.

Fig. 1 - Loose Smut



Loose Smute of Wheat



1000X

Key Points

- Grain and head replaced with loose mass of black spores.
- No smell to spores.
- Spores from smutted heads infect developing seed in healthy heads. Planting infected seed the next fall results in plants that produce loose smutted heads the next spring.


CONTROL

Loose smut and common bunt can be controlled with seed treatments (refer to 2001 OSU Extension Agents' Handbook, OCES Pub. No. E-832). An additional practice that may help control common bunt is to plant wheat early when soils are warm (>25 C or >77 F) because infection by common bunt is favored in cool soils.

Fig. 2 - Common Bunt



Healthy Head Common Bunted



1000X

Key Points

- Grain contents replaced with a dark mass of spores, but seed coat is intact.
- Spores have a rotting fishy smell.
- Spores from bunted heads survive in the soil or are carried on infested seed. These spores infect seedlings in the fall resulting in bunted heads the next spring.



Healthy Grain

Common bunted kernels

Fig. 2 - Karnal Bunt



1000X

Key Points

- Grain is filled with a dark mass of spores, but seed coat is intact and infection usually is at one end of the seed.
- Spores have a rotting, fishy smell.
- Karnal bunt spores survive in soil or are carried on seed. Infection occurs when wheat flowers, resulting in partially or fully bunted grain.

CONTROL

Control of Karnal bunt is difficult because infection occurs when wheat is flowering. Hence, seed treatments will reduce the number of spores on the seed coat, but spores in the soil may still be present at flowering when infection occurs. Karnal bunt spores can survive in soil for at least 3-5 years. Hence, rotation to non-hosts may reduce spores in the soil, but will not result in complete control in a 1-3 year period. Resistance to Karnal bunt has been reported, but the reactions of varieties adapted to Oklahoma are not known because research involving Karnal bunt can not be conducted in most areas of the United States.