



Pest e-alerts



*Entomology and Plant Pathology, Oklahoma State University
127 Noble Research Center, Stillwater, OK 74078
405.744.5527*

Vol. 16, No. 31

<http://entopl.okstate.edu/pddl/pdidl>

August 21, 2017

Early Rains May Bring Pecan Weevil Out

Phil Mulder – Oklahoma State University
Extension Entomologist and Department Head

With most of the early season insect problems behind us it is time to pay more attention to pecan weevil season. Now is the time to prepare for the upcoming barrage of weevils that can infest this season's crop. Damage from pecan weevil can vary from year to year and orchard to orchard. In fact, susceptibility to this pest can vary from tree to tree in a single, home yard situation. Reports from years past have surfaced of 95% crop loss due to pecan weevil injury. In addition to the injury on the crop itself, buyers and processors who purchase affected pecans have to either clean them out, or dispose of the product. In the native pecan growing range, this pest has evolved in nearly perfect synchrony with the host and is common throughout much of the production areas of Texas and Oklahoma. In recent years, however, the pecan weevil occasionally finds passage into the "western irrigated area." Through all the efforts of growers, state agriculture departments, and scientists, much of the New Mexico area remains weevil free. I'm sure it will be an ongoing and occasionally exhaustive effort to keep the pecan weevil from establishing a foothold in that part of the country.

The typical life cycle of the pecan weevil requires 2 to 3 years, with the majority of weevils emerging two years after burrowing into the soil. Once a weevil larva enters the soil it has likely matured through four instars. The final instar (fourth) may feed for a short while in the nut (about 1-2 weeks) and then burrows out of the nut and drops to the ground generally spending one to two years in the soil without feeding. Larvae will burrow into the soil at various depths depending on soil type, moisture, and condition of the orchard floor. Most larvae are likely 6-9 inches deep in the soil profile. Ninety percent of the larvae that entered the soil on "year 1" will pupate in "year 2." The remainder of the population will not pupate until "year 3." The pupal stage lasts for about 3 weeks (longer in females than in males) and the adults then remain inactive within the earthen cell constructed earlier by the larva and do not attempt emergence from the soil until "year 3" (about 2 years after they entered the soil as larvae). As stated previously, about 10% of the population will

not emerge until “year 4.” After emergence from the soil adult weevils live for about 2 to 3 weeks, with females surviving longer than males. Female weevils are capable of surviving much longer if nuts are not acceptable for oviposition (egg-laying). If nuts are not ready for oviposition by female weevils, then adults will cause feeding damage on the nuts. This damage may consist of feeding punctures, prior to shell hardening, which will cause premature drop of the pecan. Feeding by pecan weevil after shell hardening may cause the deterioration of the gel resulting in “sticktights.” If the weevil penetrates only the shuck, then damage may be limited to slight scars, black spots, pits or molds on the kernel near the puncture site.



Oviposition by female pecan weevils can occur as early as 2 days after emergence; however, the majority of egg production occurs 10-12 days after emergence. Regardless of how soon oviposition begins, it is initiated on early maturing varieties sooner than on trees that have late maturing nuts. Currently, the Pawnee pecans at the OSU Cimmaron Valley Research Station are in the dough stage; however, the Kanza pecans are still in the gel stage. Our first treatment will likely go out sometime this week. Each female can average around 35-75 eggs deposited during her life. She will average about four eggs per nut. With a healthy weevil population, this can account for a great amount of damage.



Much of the damage experienced by pecan weevil can be avoided with timely use of insecticides. Making the decision on when to treat; however, can be a dilemma for many growers. Although increased weevil emergence can occur after heavy rainfall events in late July and August, in Oklahoma the majority emerge in September, but this can be highly variable depending on those list of factors mentioned previously, and this year has been abnormally wet in many pecan growing regions. The decision to treat should be made with some sort of trapping program established in the

orchard. For trapping we suggest the pecan weevil Circle trap. Although growers have become familiar with these traps over the last few years, many questions remain on how to best utilize the traps. Several studies conducted in Oklahoma over the years have helped provide some answers to many of these questions. Following are several suggestions on the utility of the Circle trap and methods for arriving at a treatment threshold based on captures.

- 1) Commercial situations (several trees) – Use as many as you can afford, to represent the land area, soil types, pecan varieties, and topography in your area.
- 2) Where possible, target indicator trees – Those with a history of weevil problems, those that are earlier than most, or those located in sandy soils.
- 3) Place two traps on each targeted tree. One on the north, west or northwest side of the tree and the other just opposite. Set them at a height suitable for your operation. If cattle are going to be grazing the orchard floor, then put the traps sufficiently high on the trunk to avoid their natural curiosity. Just remember that someone needs to check traps regularly, therefore, they should be reachable for the person doing the checking. Where cattle are present, a small step ladder or truck pickup bed will help. Soon after placement on the trees, cattle lose their curiosity with these traps, probably for two reasons. First they're made of screening material, which they don't prefer to chew on and second because after a few days they tend to recognize the trap as part of the tree.



- 4) When attempting to encircle the tree trunk, place additional wire screening wherever the traps don't match up, and curl the top lip of the screening over to funnel the weevils up into the trap tops.
- 5) Mow grass under the trees to avoid providing weevils a spot to take off (fly) into the canopy.
- 6) Begin trapping by July 15 if possible, but don't be overly concerned if the nuts are still in the water stage.
- 7) Do not cease trapping too early (before mid-October).
- 8) Inspect traps at least every 2 to 3 days. Set up a M-W-F checking schedule, and stick to it.
- 9) Keep a running set of written records, to include date of trap inspection, number of weevils caught, sex of weevils caught, and tree number or location. This information should be retained for long term records and can help you identify indicator spots and/or trees.
- 10) Convert the total number of weevils collected from all traps on each inspection date into the number of weevils/trap/day. In other words, the total number of weevils from "X" traps after "X" number of days.

EXAMPLE: You examined 16 traps on 8 trees on Monday and found nothing but on Wednesday you gathered up 12 weevils from the same 16 traps. Therefore, divide 12 by (16 times 2). The two represents the days since your last check of the traps. $12/32 = 0.375$ weevils/trap/day.

- 11) Begin control measures if the number of weevils equals or exceeds 0.3 weevils/trap/day.
- 12) Be sure to regularly clean the trap screen tops to remove spiders, lacewings, webbing and dead insects which may interfere with functioning of the trap. A stiff toothbrush can help in this task.
- 13) Be diligent in this task from July to October, but really focus immediately following any 1"-2" rainfall. If conditions are hot and dry, as they often are in July and August, weevil emergence may be delayed but can still be significant up until shuck-split. In Oklahoma, populations generally peak about the third week of September.
- 14) When an application is about to be made, remove the top cylinder of each trap to allow everything to crawl through. The air-blast sprayers and the insecticide will cause a great many weevils to fall shortly after treatment. These "sick weevils" will begin crawling up tree trunks and entering traps. This can give the impression of an ineffective treatment or that you had a fresh emergence. Leave tops off for a period of 7 days, then begin the monitoring program again.

For homeowners that may have one or a few trees, it may be possible to "trap-out" the majority of their weevil population by encircling the trees with traps and following the same suggestions. This may reduce the number of insecticide applications needed in this sensitive environment and yet preserve pecans for harvesting. In localized "hot pockets," where weevils have relied on a few trees in the area, thorough monitoring and treatment over a two to three year period can dramatically reduce or even eliminate the pecan weevil for several years.

Please keep in mind that the original opening on the trap top was designed for the boll weevil, which is a much smaller weevil, so be sure to check that the opening is somewhat enlarged for the larger pecan weevil. Traps can be obtained for around \$18 each from Pecan Ag Equipment in Bristow, Oklahoma (1-918-367-5529).

Co-Editors: Eric Rebek and Justin Talley; Oklahoma Cooperative Extension Service

The pesticide information presented in this publication was current with federal and state regulations at the time of printing. The user is responsible for determining that the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label directions. The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies: Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources.